

The D -wave heavy-light mesons from QCD sum rules

Dan Zhou¹, Er-Liang Cui¹, Hua-Xing Chen^{1*},[†] Li-Sheng Geng^{1,‡}, Xiang Liu^{2,3§},[¶] and Shi-Lin Zhu^{4,5,6**},^{††}

¹*School of Physics and Nuclear Energy Engineering and International Research Center for Nuclei and Particles in the Cosmos, Beihang University, Beijing 100191, China*

²*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

³*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

⁴*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

⁵*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

⁶*Center of High Energy Physics, Peking University, Beijing 100871, China*

We study the D -wave $\bar{c}s$ heavy meson doublets $(1^-, 2^-)$ and $(2^-, 3^-)$ using the method of QCD sum rule in the framework of heavy quark effective theory. Choosing the same threshold values ω_c around 2.7 GeV, we calculate the masses of the 1^- and 3^- states. They are $m_{D_{s1}^*} = 2.81 \pm 0.10$ GeV and $m_{D_{s3}^*} = 2.85 \pm 0.08$ GeV, consistent with the newly observed $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ states by LHCb. The masses of their 2^- partners are calculated to be 2.82 ± 0.10 and 2.81 ± 0.08 GeV. The mass splittings within the same doublet are calculated to be $m_{D_{s2}} - m_{D_{s1}^*} = 0.016 \pm 0.007$ GeV and $m_{D_{s3}^*} - m_{D_{s2}} = 0.039 \pm 0.014$ GeV.

PACS numbers: 14.40.Lb, 12.38.Lg, 12.39.Hg

I. INTRODUCTION

Since the observation of $D_{s0}^*(2317)$ in 2003 [1], more and more charmed-strange mesons have been reported experimentally, which include $D_{s1}(2460)$ [2], $D_{s1}(2710)$ [3, 4], $D_{sJ}(2860)$ [4, 5], and $D_{sJ}(3040)$ [4] (see Ref. [6] for a concise review). Very recently, the LHCb Collaboration announced the observation of two charmed-strange mesons $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ with the resonance parameters [7, 8]:

$$\begin{aligned} m_{D_{s1}^*(2860)} &= (2859 \pm 12 \pm 6 \pm 23) \text{ MeV}, \\ \Gamma_{D_{s1}^*(2860)} &= (159 \pm 23 \pm 27 \pm 72) \text{ MeV}, \\ m_{D_{s3}^*(2860)} &= (2860.5 \pm 2.6 \pm 2.5 \pm 6.0) \text{ MeV}, \\ \Gamma_{D_{s3}^*(2860)} &= (53 \pm 7 \pm 4 \pm 6) \text{ MeV}. \end{aligned}$$

In addition, LHCb specified that it is the first time to identify a spin-3 resonance $D_{s3}^*(2860)$ [7, 8]. At present, the charmed-strange meson family is becoming more and more abundant with the experimental progress.

Until now, there are good candidates of the $1S$ and $1P$ states in the charmed-strange meson family [9]. These newly observed charmed-strange mesons provide a good platform to study the properties of the higher radial and orbital excitations of the charmed-strange meson. For example, in Ref. [10] Sun and Liu suggested that $D_{sJ}(3040)$ can be a good candidate of the $2P$ state in the charmed-strange meson family, which is the radial excitation of $D_{s1}(2460)$. The recently reported $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ states stimulated extensive discussions of whether they can be categorized into the $1D$

charmed-strange mesons [11–13]. In Ref. [11], the two-body strong decays of $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ as the 1^3D_1 and 1^3D_3 states in charmed-strange meson family were studied by the quark pair creation model, which shows that $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ are the 1^3D_1 and 1^3D_3 states, respectively. Later, Wang studied $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ using the effective Lagrangian approach [12]. Recently, Godfrey and Moats [13] indicated that $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ are the 1^3D_1 and 1^3D_3 charmed-strange mesons, respectively. Thus, the results in Refs. [12, 13] supports the assignment of $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ proposed in Ref. [11].

In this paper we shall use the method of QCD sum rule to study the D -wave heavy meson doublets $(1^-, 2^-)$ and $(2^-, 3^-)$ containing one heavy anti-quark and one strange quark [14, 15]. We shall work in the framework of the heavy quark effective theory (HQET) [16–18], which has been successful to study heavy hadrons containing a single heavy quark. The mass of the ground state heavy mesons was studied in Refs. [19–24]. The masses of the lowest excited non-strange heavy meson doublets $(0^+, 1^+)$ and $(1^+, 2^+)$ were studied in Refs. [26–28]. The masses of the lowest excited $\bar{c}s$ heavy mesons in the $(0^+, 1^+)$ and $(1^+, 2^+)$ doublets were studied in Ref. [29]. There were also some early studies using the method of QCD sum rules but in full QCD [30, 31]. In this paper we shall follow the procedures used in Refs. [25–27, 29], and study the D -wave $\bar{c}s$ heavy meson in the $(1^-, 2^-)$ and $(2^-, 3^-)$ doublets. We shall also follow Refs. [25–27, 29] and consider the $O(1/m_Q)$ corrections, where m_Q is the heavy quark mass.

This paper is organized as follows. In Sec. II, we introduce the interpolating currents for the D -wave $\bar{c}s$ heavy meson doublets $(1^-, 2^-)$ and $(2^-, 3^-)$, and use them to perform QCD sum rule analyses at the leading order. Then in Sec. III we calculate the $O(1/m_Q)$ corrections. The results are summarized and discussed in Sec. IV.

*Corresponding author

§Corresponding author

**Corresponding author

[†]Electronic address: hxchen@buaa.edu.cn

[‡]Electronic address: lisheng.geng@buaa.edu.cn

[¶]Electronic address: xiangliu@lzu.edu.cn

^{††}Electronic address: zhushl@pku.edu.cn

II. THE SUM RULES AT THE LEADING ORDER (IN THE $m_Q \rightarrow \infty$ LIMIT)

The interpolating currents for the heavy mesons with arbitrary spin and parity have been studied and given in Refs. [25–27]. Here we briefly discuss how we obtain the interpolating currents coupling to the D -wave $\bar{c}s$ heavy meson doublets ($1^-, 2^-$) and ($2^-, 3^-$). We use J_{j,P,j_l} to denote these fields where the first two subscripts denote the spin and parity of the heavy mesons, and the last subscript denotes the angular momentum of the light components. We also use the following notations: $\gamma_t^\mu = \gamma^\mu - \not{v}v^\mu$, $D_t^\mu = D^\mu - (D \cdot v)v^\mu$, with $D^\mu = \partial^\mu - igA^\mu$ is the gauge-covariant derivative, $\bar{h}_v$ is the heavy quark field in HQET, v is the velocity of the heavy quark, and $g_t^{\alpha_1\alpha_2} = g^{\alpha_1\alpha_2} - v^{\alpha_1}v^{\alpha_2}$ is the transverse metric tensor.

Based on the pseudoscalar current $\bar{h}_v\gamma_5 q$ of $J^P = 0^-$ and the vector current $\bar{h}_v\gamma_\mu q$ of $J^P = 1^-$, we can construct the D -wave interpolating currents, by adding two extra derivatives. Assuming these two derivatives are both acting on the light quark, it can have either $j_l^{P_l} = 3/2^+$:

$$\mathcal{D}_t^\alpha \mathcal{D}_t^\beta \times \gamma_\beta \gamma_5 q, \quad (1)$$

or $j_l^{P_l} = 5/2^+$:

$$\mathcal{D}_t^{\alpha_1} \mathcal{D}_t^{\alpha_2} \times q. \quad (2)$$

Use the light quark of $j_l^{P_l} = 3/2^+$, we can construct the interpolating currents coupling to the D -wave ($1^-, 2^-$) spin doublet. One of them has the pseudoscalar structure, while the other has the vector structure:

$$J_{x,-,3/2}^{\dagger\alpha} = \bar{h}_v \gamma_5 \times \mathcal{D}_t^\alpha \mathcal{D}_t^\beta \times \gamma_\beta \gamma_5 q, \quad (3)$$

$$J_{y,-,3/2}^{\dagger\alpha_1\alpha_2} = \bar{h}_v \gamma_t^{\alpha_2} \times \mathcal{D}_t^{\alpha_1} \mathcal{D}_t^{\alpha_2} \times \gamma_\beta \gamma_5 q. \quad (4)$$

However, they are not pure 1^- or 2^- (Particularly, $J_{y,-,3/2}^{\dagger\alpha_1\alpha_2}$ contains both 1^- and 2^- components). Therefore, we need to do an extra process to project out their 1^- and 2^- components:

$$J_{1,-,3/2}^{\dagger\alpha} = \sqrt{\frac{3}{4}} \bar{h}_v (-i) (\mathcal{D}_t^\alpha - \frac{1}{3} \gamma_t^\alpha \not{D}_t) \not{D}_t q, \quad (5)$$

$$J_{2,-,3/2}^{\dagger\alpha_1\alpha_2} = \sqrt{\frac{1}{2}} \bar{h}_v \gamma_5^5 \frac{(-i)^2}{2} (\gamma_t^{\alpha_1} \mathcal{D}_t^{\alpha_2} \not{D}_t + \gamma_t^{\alpha_2} \mathcal{D}_t^{\alpha_1} \not{D}_t - \frac{2}{3} g_t^{\alpha_1\alpha_2} \mathcal{D}_t \cdot \mathcal{D}_t) q, \quad (6)$$

where we have modified their expressions to be consistent with Refs. [25–27]. Similarly, we can construct the following interpolating currents to study the D -wave ($2^-, 3^-$) spin doublet:

$$J_{2,-,5/2}^{\dagger\alpha_1\alpha_2} = \sqrt{\frac{5}{6}} \bar{h}_v \gamma_5^5 \frac{(-i)^2}{2} (\mathcal{D}_t^{\alpha_2} \mathcal{D}_t^{\alpha_1} + \mathcal{D}_t^{\alpha_1} \mathcal{D}_t^{\alpha_2} - \frac{2}{5} \mathcal{D}_t^{\alpha_2} \gamma_t^{\alpha_1} \not{D}_t - \frac{2}{5} \mathcal{D}_t^{\alpha_1} \gamma_t^{\alpha_2} \not{D}_t - \frac{2}{5} g_t^{\alpha_1\alpha_2} \mathcal{D}_t \cdot \mathcal{D}_t) q, \quad (7)$$

$$J_{3,-,5/2}^{\dagger\alpha_1\alpha_2\alpha_3} = \sqrt{\frac{1}{2}} \bar{h}_v S_1 [\gamma_t^{\alpha_1} (-i)^2 \mathcal{D}_t^{\alpha_2} \mathcal{D}_t^{\alpha_3}] q, \quad (8)$$

where S_1 denotes symmetrization and subtracting the trace terms in the sets $(\alpha_1\alpha_2\alpha_3)$.

Based on the scalar current $\bar{h}_v q$ of $J^P = 0^+$ and the axial-vector current $\bar{h}_v \gamma_\mu \gamma_5 q$ of $J^P = 1^+$, we can also construct the D -wave interpolating currents, by adding only one derivative. For example, the light quark containing one derivative can have $j_l^{P_l} = 3/2^-$:

$$\mathcal{D}_t^\alpha \times q. \quad (9)$$

Then it can be used to construct the interpolating currents coupling to the D -wave ($1^-, 2^-$) spin doublet:

$$J_{x,-,3/2}^{\dagger\alpha} = \bar{h}_v \times \mathcal{D}_t^\alpha \times q, \quad (10)$$

$$J_{y,-,3/2}^{\dagger\alpha_1\alpha_2} = \bar{h}_v \gamma_t^{\alpha_2} \gamma_5 \times \mathcal{D}_t^{\alpha_1} \times q. \quad (11)$$

These two currents can be further manipulated to obtain the pure 1^- and 2^- currents.

$$J_{1,-,3/2}^{\dagger\alpha} = \sqrt{\frac{3}{4}} \bar{h}_v (-i) (\mathcal{D}_t^\alpha - \frac{1}{3} \gamma_t^\alpha \not{D}_t) q, \quad (12)$$

$$J_{2,-,3/2}^{\dagger\alpha_1\alpha_2} = \sqrt{\frac{1}{2}} \bar{h}_v \gamma_5^5 \frac{(-i)^2}{2} (\gamma_t^{\alpha_1} \mathcal{D}_t^{\alpha_2} + \gamma_t^{\alpha_2} \mathcal{D}_t^{\alpha_1} - \frac{2}{3} g_t^{\alpha_1\alpha_2} \not{D}_t) q. \quad (13)$$

However, compared with these currents containing one derivative, the currents containing two derivatives seem to describe the internal structure of the D -wave heavy mesons in a more appropriate way, so we shall use Eqs. (5), (6), (7) and (8) to perform QCD sum rule analyses. Moreover, the two currents $J_{1,-,3/2}$ and $J_{2,-,3/2}$ give identical sum rules at the leading order, and the sum rules at the $O(1/m_Q)$ order can be obtained using either of them [25–27, 29] (ideally the results should be identical, while actually they have small differences but negligible). Accordingly, we only need to use one of them to perform QCD sum rule analyses. So do $J_{2,-,5/2}$ and $J_{3,-,5/2}$. In the following discussions we shall use $J_{1,-,3/2}$ and $J_{3,-,5/2}$, because they couple to the newly observed $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ states by LHCb [7, 8].

In the $m_Q \rightarrow \infty$ limit we can assume $|j, P, j_l\rangle$ to be the heavy meson state with the quantum numbers j, P and j_l , and the relation between this state and the relevant interpolating field is

$$\langle 0 | J_{j,P,j_l}^{\alpha_1 \dots \alpha_j} | j', P', j_l' \rangle = f_{P,j_l} \delta_{jj'} \delta_{PP'} \delta_{j_l j_l'} \eta_t^{\alpha_1 \dots \alpha_j}, \quad (14)$$

where f_{P,j_l} is the decay constant. It has the same value for the two states in the same doublet in the $m_Q \rightarrow \infty$ limit. $\eta_t^{\alpha_1 \dots \alpha_j}$ is the transverse, symmetric, and traceless polarization tensor. In this paper we need to use η_t^α and $\eta_t^{\alpha_1\alpha_2\alpha_3}$ which have the following property at the leading order

$$\eta_t^\alpha \eta_t^{\beta*} = \tilde{g}_t^{\alpha\beta}, \quad (15)$$

$$\eta_t^{\alpha_1\alpha_2\alpha_3} \eta_t^{\beta_1\beta_2\beta_3*} = S_2 [\tilde{g}_t^{\alpha_1\beta_1} \tilde{g}_t^{\alpha_2\beta_2} \tilde{g}_t^{\alpha_3\beta_3}], \quad (16)$$

where $\tilde{g}^{\mu\nu} = g^{\mu\nu} - q^\mu q^\nu / m^2$ and S_2 denotes symmetrization and subtracting the trace terms in the sets $(\alpha_1\alpha_2\alpha_3)$ and $(\beta_1\beta_2\beta_3)$.

Using the two interpolating currents $J_{1,-,3/2}$ and $J_{3,-,5/2}$, we can construct the two-point correlation function

$$\begin{aligned}\Pi_{j,P,j_l}^{\alpha_1 \dots \alpha_j \beta_1 \dots \beta_j}(\omega) &= i \int d^4x e^{ikx} \langle 0 | T [J_{j,P,j_l}^{\alpha_1 \dots \alpha_j}(x) J_{j,P,j_l}^{\dagger \beta_1 \dots \beta_j}(0)] | 0 \rangle \\ &= (-1)^j S_3 [g_t^{\alpha_1 \beta_1} \dots g_t^{\alpha_j \beta_j}] \Pi_{j,P,j_l}(\omega),\end{aligned}\quad (17)$$

where $\omega = 2v \cdot k$ is twice the external off-shell energy, and S_3 denotes symmetrization and subtracting the trace terms in the sets $(\alpha_1 \dots \alpha_j)$ and $(\beta_1 \dots \beta_j)$. At the hadron level, it can be written as

$$\Pi_{j,P,j_l}(\omega) = \frac{f_{P,j_l}^2}{2\bar{\Lambda}_{j,P,j_l} - \omega} + \text{higher states}, \quad (18)$$

where $\bar{\Lambda}_{j,P,j_l} = \lim_{m_Q \rightarrow \infty} (m_{j,P,j_l} - m_Q)$, and m_{j,P,j_l} is the mass of the lowest-lying heavy meson state which $J_{j,P,j_l}^{\alpha_1 \dots \alpha_j}(x)$ couples to. At the quark and gluon level, we can calculate the two-point correlation function (17) using the method of QCD sum rule. To do this we follow the approaches used in Refs. [25–27, 29]. After inserting Eq. (5) and (8) into Eq. (17), and performing the Borel transformation, we obtain

$$\begin{aligned}\Pi_{1,-,3/2}(\omega_c, T) &= f_{-,3/2}^2 e^{-2\bar{\Lambda}_{1,-,3/2}/T} \\ &= \frac{7}{2560\pi^2} \int_{2m_s}^{\omega_c} [\omega^6 + 2m_s\omega^5 - 10m_s^2\omega^4] e^{-\omega/T} d\omega \\ &\quad - \frac{1}{8\pi} \langle \alpha_s GG \rangle T^3,\end{aligned}\quad (19)$$

$$\begin{aligned}\Pi_{3,-,5/2}(\omega_c, T) &= f_{-,5/2}^2 e^{-2\bar{\Lambda}_{3,-,5/2}/T} \\ &= \frac{1}{640\pi^2} \int_{2m_s}^{\omega_c} [\omega^6 + 2m_s\omega^5 - 10m_s^2\omega^4] e^{-\omega/T} d\omega \\ &\quad - \frac{3}{32\pi} \langle \alpha_s GG \rangle T^3.\end{aligned}\quad (20)$$

We note that there are $2 \times 2 = 4$ derivatives, and so the calculations are not easy. To deal with them, we have used a software called *Mathematica* with a package called *FeynCalc* [32]. Moreover, we do not consider the radiative corrections in order to simplify our calculations.

Particularly, the quark condensate $\langle \bar{q}q \rangle$ and the mixed condensate $\langle g_s \bar{q} \sigma G q \rangle$ both vanish in this case. This is much different from those sum rules for $(0^+, 1^+)$ and $(1^+, 2^+)$ doublets [25–27, 29], and it makes the convergence of Eq. (19) and (20) very good. To clearly see this, we show the convergence of Eq. (20) in Fig. 1, where ω_c is taken to be 2.7 GeV, and the following values for the gluon condensate and the strange quark mass are used [25–27, 29, 33]:

$$\langle \frac{\alpha_s}{\pi} GG \rangle = 0.005 \pm 0.004 \text{ GeV}^4, \quad (21)$$

$$m_s = 0.15 \text{ GeV}. \quad (22)$$

Finally, we differentiate Eq. (19) and (20) with respect to $-2/T$, divide the results by themselves, and obtain

$$\Lambda_{j,P,j_l}(\omega_c, T) = \frac{\frac{\partial}{\partial(-2/T)} \Pi_{j,P,j_l}(\omega_c, T)}{\Pi_{j,P,j_l}(\omega_c, T)}. \quad (23)$$




FIG. 1: Various contributions to Eq. (20), as functions of the Borel mass M_B in units of GeV^7 at $\omega_c = 2.7 \text{ GeV}$. The labels indicate the dimension up to which the OPE terms are included.

The results can be further used to evaluate f_{P,j_l} :

$$f_{P,j_l}(\omega_c, T) = \sqrt{\Pi_{j,P,j_l}(\omega_c, T) \times e^{2\bar{\Lambda}_{j,P,j_l}(\omega_c, T)/T}}. \quad (24)$$

Here we note again that the sum rule obtained by using the other current $J_{2,-,3/2}$ is very similar to Eq. (19). Hence, we have $\Lambda_{1,-,3/2} = \Lambda_{2,-,3/2}$, and we shall use another symbol $\Lambda_{-,3/2}$ to denote them. Similarly we shall use the symbol $\Lambda_{-,5/2}$ to denote $\Lambda_{2,-,5/2}$ and $\Lambda_{3,-,5/2}$. To differentiate the masses within the same doublet, we need to work at the $O(1/m_Q)$ order, which will be done in the next section.

To perform the numerical analysis, firstly we require that the high-order power corrections be less than 30% of the perturbation term, and obtain the minimum value T_{min} of the allowed Borel parameter; secondly we require that the pole contribution

$$\text{Pole contribution} \equiv \frac{\Pi_{j,P,j_l}(\omega_c, T)}{\Pi_{j,P,j_l}(\infty, T)}, \quad (25)$$

is larger than 30%, and obtain the maximum value T_{max} of the allowed Borel parameter. Altogether we have the working interval $T_{min} < T < T_{max}$ for a fixed ω_c . In the sum rules (19) and (20) ω_c is free parameter, and we choose it to be around 2.7 GeV for both $\Pi_{1,-,3/2}$ and $\Pi_{3,-,5/2}$, because the $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ observed by LHCb have similar masses 2859 MeV and 2860.5 MeV [7]. Using this $\omega_c = 2.7 \text{ GeV}$, we obtain our working regions, around $0.35 \text{ GeV} < T < 0.48 \text{ GeV}$ for $J_{1,-,3/2}$, and around $0.39 \text{ GeV} < T < 0.47 \text{ GeV}$ for $J_{3,-,5/2}$. As an example, we show the comparison between the pole and continuum contributions for $J_{3,-,5/2}$ in Fig. 2.

Finally, we solve Eq. (23) and Eq. (24), and evaluate Λ_{-,j_l} and f_{-,j_l} . We show the variations of $\Lambda_{-,3/2}$ and $f_{-,3/2}$ with respect to the Borel mass T and the threshold value ω_c in Fig. 3. These figures are shown in the region $0.25 \text{ GeV} < T < 0.55 \text{ GeV}$, but we find that their dependence on the Borel mass T becomes weaker in our working region $0.35 \text{ GeV} < T < 0.48 \text{ GeV}$. We obtain the following numerical results:

$$\Lambda_{-,3/2} = 1.10 \pm 0.06 \text{ GeV}, \quad (26)$$

$$f_{-,3/2} = 0.19 \pm 0.05 \text{ GeV}^{7/2}, \quad (27)$$




FIG. 2: The solid curve shows the pole contribution and the dashed curve shows the continuum contribution (= 1-pole contribution), when $J_{3,-5/2}$ is used and ω_c is fixed to be 2.7 GeV.

where the central value corresponds to $T = 0.42$ GeV and $\omega_c = 2.7$ GeV.




FIG. 3: The variation of $\Lambda_{-3/2}$ and $f_{-3/2}$ with respect to the Borel mass T and the threshold value ω_c . The short-dashed, solid and long-dashed curves are obtained by fixing $\omega_c = 2.5, 2.7$ and 2.9 GeV, respectively. Our working region is $0.35 \text{ GeV} < T < 0.48 \text{ GeV}$.

Similarly, we show the variations of $\Lambda_{-5/2}$ and $f_{-5/2}$ in Fig. 4. These figures are shown in the region $0.30 \text{ GeV} < T < 0.60 \text{ GeV}$. Again we find that their dependence on the Borel mass T becomes weaker in our working region $0.39 \text{ GeV} < T < 0.47 \text{ GeV}$. We obtain the following numerical

results:

$$\Lambda_{-5/2} = 1.14 \pm 0.05 \text{ GeV}, \quad (28)$$

$$f_{-5/2} = 0.15 \pm 0.04 \text{ GeV}^{7/2}, \quad (29)$$

where the central value corresponds to $T = 0.43 \text{ GeV}$ and $\omega_c = 2.7 \text{ GeV}$.




FIG. 4: The variation of $\Lambda_{-5/2}$ and $f_{-5/2}$ with respect to the Borel mass T and the threshold value ω_c . The short-dashed, solid and long-dashed curves are obtained by fixing $\omega_c = 2.5, 2.7$ and 2.9 GeV, respectively. Our working region is $0.39 \text{ GeV} < T < 0.47 \text{ GeV}$.

III. THE SUM RULES AT THE $O(1/m_Q)$ ORDER

The Lagrangian of HQET, up to the $O(1/m_Q)$ order, can be written as [27, 29]

$$\mathcal{L}_{\text{eff}} = \bar{h}_v i v \cdot D h_v + \frac{1}{2m_Q} \mathcal{K} + \frac{1}{2m_Q} \mathcal{S}, \quad (30)$$

where $\mathcal{K}$ is the operator of the nonrelativistic kinetic energy with a negative sign:

$$\mathcal{K} = \bar{h}_v (i D_\perp)^2 h_v, \quad (31)$$

and $\mathcal{S}$ is the Pauli term to describe the chromomagnetic interaction:

$$\mathcal{S} = \frac{g}{2} C_{\text{mag}}(m_Q/\mu) \bar{h}_v \sigma_{\mu\nu} G^{\mu\nu} h_v, \quad (32)$$

where $C_{\text{mag}}(m_Q/\mu) = [\alpha_s(m_Q)/\alpha_s(\mu)]^{3/\beta_0}$ and $\beta_0 = 11 - 2n_f/3$.

We use δm and δf to denote the corrections to the mass m_{j,P,j_l} and the coupling constant f_{P,j_l} at the $O(1/m_Q)$ order. The pole term on the hadron side, Eq. (18), can be written as:

$$\begin{aligned}\Pi(\omega)_{pole} &= \frac{(f + \delta f)^2}{2(\bar{\Lambda} + \delta m) - \omega} \\ &= \frac{f^2}{2\bar{\Lambda} - \omega} - \frac{2\delta m f^2}{(2\bar{\Lambda} - \omega)^2} + \frac{2f\delta f}{2\bar{\Lambda} - \omega}.\end{aligned}\quad (33)$$

In this paper we shall only evaluate δm . To do this, we use the Lagrangian (30) defined at the $O(1/m_Q)$ order, and consider the following three-point correlation functions

$$\begin{aligned}\delta_O \Pi_{j,P,i}^{\alpha_1 \dots \alpha_j, \beta_1 \dots \beta_j}(\omega, \omega') \\ = i^2 \int d^4x d^4y e^{ik \cdot x - ik' \cdot y} \langle 0 | T [J_{j,P,i}^{\alpha_1 \dots \alpha_j}(x) O(0) J_{j,P,i}^{\beta_1 \dots \beta_j}(y)] | 0 \rangle \\ = (-1)^j S_2 [g_t^{\alpha_1 \beta_1} \dots g_t^{\alpha_j \beta_j}] \delta_O \Pi_{j,P,j_l}(\omega),\end{aligned}\quad (34)$$

where $O = \mathcal{K}$ or $\mathcal{S}$. At the hadron level, we can pick their pole parts

$$\begin{aligned}\delta_{\mathcal{K}} \Pi(\omega, \omega')_{j,P,j_l} &= \frac{f^2 K_{P,j_l}}{(2\bar{\Lambda} - \omega)(2\bar{\Lambda} - \omega')} + \frac{f^2 G_{\mathcal{K}}(\omega')}{2\bar{\Lambda} - \omega} \\ &\quad + \frac{f^2 G_{\mathcal{K}}(\omega)}{2\bar{\Lambda} - \omega'},\end{aligned}\quad (35)$$

$$\begin{aligned}\delta_{\mathcal{S}} \Pi(\omega, \omega')_{j,P,j_l} &= \frac{d_M f^2 \Sigma_{P,j_l}}{(2\bar{\Lambda} - \omega)(2\bar{\Lambda} - \omega')} + \frac{d_M f^2 G_{\mathcal{S}}(\omega')}{2\bar{\Lambda} - \omega} \\ &\quad + \frac{d_M f^2 G_{\mathcal{S}}(\omega)}{2\bar{\Lambda} - \omega'},\end{aligned}\quad (36)$$

where

$$\begin{aligned}K_{P,j_l} &= \langle j, P, j_l | \bar{h}_v (iD_\perp)^2 h_v | j, P, j_l \rangle, \\ 2d_M \Sigma_{P,j_l} &= \langle j, P, j_l | g \bar{h}_v \sigma_{\mu\nu} G^{\mu\nu} h_v | j, P, j_l \rangle, \\ d_M &= d_{j,j_l}, \\ d_{j_l-1/2,j_l} &= 2j_l + 2 \\ d_{j_l+1/2,j_l} &= -2j_l.\end{aligned}\quad (37)$$

From these equations we know that the term $\mathcal{S}$ causes a mass splitting within the same doublet, while the term $\mathcal{K}$ does not. Moreover, the term $\mathcal{S}$ can also cause a mixing of states with the same j, P but different j_l , such as a mass splitting between $|2, -, 3/2\rangle$ and $|2, -, 5/2\rangle$. This effect has been studied in Ref. [34], where its corrections are found to be negligible. Hence, we do not consider this effect in this paper.

Fixing $\omega = \omega'$ and comparing Eq. (33), Eq. (35) and Eq. (36), we obtain

$$\delta m_{P,j_l} = -\frac{1}{4m_Q} (K_{P,j_l} + d_M C_{mag} \Sigma_{P,j_l}). \quad (38)$$

At the quark and gluon level, we can calculate Eqs. (34) using the method of QCD sum rule, and evaluate K_{P,j_l} and Σ_{P,j_l} . To do this, again we follow the approaches used in Refs. [27, 29]: after inserting Eq. (5) and (8) into Eq. (34), we

make a double Borel transformation for both ω and ω' , and obtain two Borel parameters T_1 and T_2 . Then we take these two Borel parameters to be equal, and obtain the following two sum rules for $K_{-,3/2}$ and $\Sigma_{-,3/2}$:

$$\begin{aligned}f_{-,3/2}^2 K_{-,3/2} e^{-2\bar{\Lambda}_{-,3/2}/T} \\ = -\frac{11}{7168\pi^2} \int_{2m_s}^{\omega_c} \omega^8 e^{-\omega/T} d\omega + \frac{91}{64\pi} \langle \alpha_s GG \rangle T^5,\end{aligned}\quad (39)$$

$$f_{-,3/2}^2 \Sigma_{-,3/2} e^{-2\bar{\Lambda}_{-,3/2}/T} = \frac{7}{240\pi} \langle \alpha_s GG \rangle T^5, \quad (40)$$

and the following two sum rules for $K_{-,5/2}$ and $\Sigma_{-,5/2}$:

$$\begin{aligned}f_{-,5/2}^2 K_{-,5/2} e^{-2\bar{\Lambda}_{-,5/2}/T} \\ = -\frac{1}{1280\pi^2} \int_{2m_s}^{\omega_c} \omega^8 e^{-\omega/T} d\omega + \frac{71}{96\pi} \langle \alpha_s GG \rangle T^5,\end{aligned}\quad (41)$$

$$f_{-,5/2}^2 \Sigma_{-,5/2} e^{-2\bar{\Lambda}_{-,5/2}/T} = \frac{1}{40\pi} \langle \alpha_s GG \rangle T^5. \quad (42)$$

We note that in these sum rules the m_s corrections are neglected.

Finally, we obtain $K_{-,3/2}$ and $\Sigma_{-,3/2}$ by simply dividing Eq. (39) and (40) by the sum rule (19), and $K_{-,5/2}$ and $\Sigma_{-,5/2}$ by simply dividing Eq. (41) and (42) by the sum rule (20). We show the variations of $K_{-,3/2}$ and $\Sigma_{-,3/2}$ with respect to the Borel mass T and the threshold value ω_c in Fig. 5 in the region $0.25 \text{ GeV} < T < 0.55 \text{ GeV}$, and their dependence on the Borel mass T becomes weaker in our working region $0.35 \text{ GeV} < T < 0.48 \text{ GeV}$. We obtain the following numerical results:

$$K_{-,3/2} = -2.25 \pm 0.36 \text{ GeV}^2, \quad (43)$$

$$\Sigma_{-,3/2} = 0.010 \pm 0.004 \text{ GeV}^2, \quad (44)$$

where the central value corresponds to $T = 0.42 \text{ GeV}$ and $\omega_c = 2.7 \text{ GeV}$.

Similarly we show the variations of $K_{-,5/2}$ and $\Sigma_{-,5/2}$ in Fig. 6 in the region $0.30 \text{ GeV} < T < 0.60 \text{ GeV}$, and their dependence on the Borel mass T becomes weaker in our working region $0.39 \text{ GeV} < T < 0.47 \text{ GeV}$. We obtain the following numerical results:

$$K_{-,5/2} = -2.16 \pm 0.28 \text{ GeV}^2, \quad (45)$$

$$\Sigma_{-,5/2} = 0.017 \pm 0.006 \text{ GeV}^2, \quad (46)$$

where the central value corresponds to $T = 0.43 \text{ GeV}$ and $\omega_c = 2.7 \text{ GeV}$.

IV. NUMERICAL RESULTS AND DISCUSSIONS

Combining the results obtained in Sec. II and Sec. III, we arrive at the following weighted average mass for the D -wave $c\bar{s}$ heavy meson doublet ($1^-, 2^-$):

$$\begin{aligned}\frac{1}{8} (3m_{D_{s1}^*} + 5m_{D_{s2}^*}) &= m_c + (1.10 \pm 0.06) \text{ GeV} \\ &\quad + \frac{1}{m_c} [(0.56 \pm 0.09) \text{ GeV}^2],\end{aligned}\quad (47)$$




FIG. 5: The variation of $K_{-3/2}$ and $\Sigma_{-3/2}$ with respect to the Borel mass T and the threshold value ω_c . The short-dashed, solid and long-dashed curves are obtained by fixing $\omega_c = 2.5, 2.7$ and 2.9 GeV, respectively. Our working region is $0.35 \text{ GeV} < T < 0.48 \text{ GeV}$.

where D_{s2} is used to denote the 2^- partner of D_{s1}^* . Their mass splitting is:

$$m_{D_{s2}} - m_{D_{s1}^*} = \frac{1}{m_c} [(0.021 \pm 0.008) \text{ GeV}^2]. \quad (48)$$

From these values, we find that the $\mathcal{O}(1/m_Q)$ corrections are important and can not be neglected.

To obtain numerical results, we use the PDG value $m_c = 1.275 \pm 0.025 \text{ GeV}$ [35] for the charm quark mass in the $\overline{\text{MS}}$ scheme. We note that one may also use its pole mass, but then the threshold value ω_c should be properly fine-tuned. Therefore, our results for the masses of the heavy mesons have large theoretical uncertainties. However, their differences within the same doublet do not depend much on the charm quark mass and the threshold value, so they are produced quite well, with much less theoretical uncertainties:

$$\begin{aligned} m_{D_{s1}^*} &= 2.81 \pm 0.10 \text{ GeV}, \\ m_{D_{s2}} &= 2.82 \pm 0.10 \text{ GeV}, \\ m_{D_{s2}} - m_{D_{s1}^*} &= 0.016 \pm 0.007 \text{ GeV}. \end{aligned} \quad (49)$$

The mass of 1^- state is $2.81 \pm 0.10 \text{ GeV}$, consistent with the D_{s1}^* (2860) newly observed by LHCb, $m_{D_{s1}^*}^{\text{exp}} = 2859 \pm 12 \pm 6 \pm 23 \text{ MeV}$ [7].

Similarly, we obtain the following weighted average mass




FIG. 6: The variation of $K_{-5/2}$ and $\Sigma_{-5/2}$ with respect to the Borel mass T and the threshold value ω_c . The short-dashed, solid and long-dashed curves are obtained by fixing $\omega_c = 2.5, 2.7$ and 2.9 GeV, respectively. Our working region is $0.39 \text{ GeV} < T < 0.47 \text{ GeV}$.

for the D -wave $\bar{c}s$ ($2^-, 3^-$) heavy meson doublet:

$$\begin{aligned} \frac{1}{12}(5m_{D'_{s2}} + 7m_{D_{s3}^*}) &= m_c + (1.14 \pm 0.05) \text{ GeV} \\ &+ \frac{1}{m_c} [(0.54 \pm 0.07) \text{ GeV}^2], \end{aligned} \quad (50)$$

where D'_{s2} is used to denote the 2^- partner of D_{s3}^* . Their mass splitting is:

$$m_{D_{s3}^*} - m_{D'_{s2}} = \frac{1}{m_c} [(0.050 \pm 0.018) \text{ GeV}^2]. \quad (51)$$

Again we find that the $\mathcal{O}(1/m_Q)$ corrections are important and have large uncertainties. Our results are:

$$\begin{aligned} m_{D'_{s2}} &= 2.81 \pm 0.08 \text{ GeV}, \\ m_{D_{s3}^*} &= 2.85 \pm 0.08 \text{ GeV}, \\ m_{D_{s3}^*} - m_{D'_{s2}} &= 0.039 \pm 0.014 \text{ GeV}. \end{aligned} \quad (52)$$

The mass of the 3^- state is $2.85 \pm 0.08 \text{ GeV}$, also consistent with the D_{s3}^* (2860) newly observed by LHCb, $m_{D_{s3}^*}^{\text{exp}} = 2860.5 \pm 2.6 \pm 2.5 \pm 6.0 \text{ MeV}$ [7].

The bs system can be similarly studied by replacing m_c by m_b and multiplying Σ_{-j_l} by $C_{\text{mag}} \approx 0.8$ [27, 29]. Here we only give their mass differences within the same doublet because their mass depends much on the bottom quark mass m_b , whose

value has large uncertainties. Using the same threshold values ω_c around 3.3 GeV and assuming $4 \text{ GeV} < m_b < 5 \text{ GeV}$, we obtain the mass differences within the same doublet

$$\begin{aligned} m_{B_{s2}} - m_{B_{s1}^*} &= 0.004 \pm 0.002 \text{ GeV}, \\ m_{B_{s3}^*} - m_{B_{s2}'} &= 0.009 \pm 0.004 \text{ GeV}. \end{aligned} \quad (53)$$

We can similarly replace the strange quark by up and down quarks and extract the masses of the non-strange D -wave heavy mesons. To do this we use slightly smaller threshold values $\omega_c \sim 2.5 \text{ GeV}$, and obtain the working region $0.39 \text{ GeV} < T < 0.43 \text{ GeV}$ for $(1^-, 2^-)$ doublet. However, there is no stability window for $(2^-, 3^-)$ doublet, unless we require the pole contribution to be greater than 20% only, and now the working region is $0.46 \text{ GeV} < T < 0.49 \text{ GeV}$. The numerical results are

$$\begin{aligned} m_{D_1^*} &= 2.75 \pm 0.09 \text{ GeV}, \\ m_{D_2} &= 2.78 \pm 0.09 \text{ GeV}, \\ m_{D_2} - m_{D_1^*} &= 0.02 \pm 0.01 \text{ GeV}, \\ m_{D_{s2}'} &= 2.72 \pm 0.10 \text{ GeV}, \\ m_{D_{s3}^*} &= 2.78 \pm 0.10 \text{ GeV}, \\ m_{D_{s3}^*} - m_{D_{s2}'} &= 0.06 \pm 0.03 \text{ GeV}. \end{aligned} \quad (54)$$

Again we note that the masses have large uncertainties, but their differences within the same doublet are produced quite well.

In summary, we have studied the D -wave $(1^-, 2^-)$ and $(2^-, 3^-)$ $\bar{c}s$ heavy meson doublets and calculated their masses up to the $O(1/m_Q)$ order using the method of QCD sum rule in the framework of HQET. The masses of 1^- and 3^- states are calculated to be $m_{D_{s1}^*} = 2.81 \pm 0.10 \text{ GeV}$ and $m_{D_{s3}^*} = 2.85 \pm 0.08 \text{ GeV}$, consistent with the newly observed $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ states by LHCb [7]. In our calculations we have chosen the same threshold value $\omega \approx 2.7 \text{ GeV}$ for both of them, and obtained a mass difference between D_{s1}^* and D_{s3}^* to be 0.04 GeV. Considering the mass uncertainties are about 0.1 GeV, our results are consistent with the experimental data [7]. The masses of their 2^- partners are calculated to be 2.82 ± 0.10 and $2.81 \pm 0.08 \text{ GeV}$. We note that our results for the masses of the heavy mesons have large theoretical uncertainties. One of its sources is the uncertainty of the charm quark mass. Besides this, we do not

consider radiative corrections in the OPE calculations, which may give extra uncertainties. However, the mass splittings within the same doublet do not depend much on this, and are reproduced quite well, i.e., $m_{D_{s2}} - m_{D_{s1}^*} = 0.016 \pm 0.007 \text{ GeV}$ and $m_{D_{s3}^*} - m_{D_{s2}'} = 0.039 \pm 0.014 \text{ GeV}$. We have also estimated their decay constants at the leading order (in the $m_Q \rightarrow \infty$ limit), that is $f_{-,3/2} = 0.19 \pm 0.05 \text{ GeV}^{7/2}$ and $f_{-,5/2} = 0.15 \pm 0.04 \text{ GeV}^{7/2}$.

At present, the two 2^- charmed-strange mesons are still missing. The predicted masses of these two 2^- charmed-strange mesons in this work can be further tested by future experiments. We also expect more experimental progresses on higher radial and orbital excitations in the charmed-strange meson family. Besides the experiments, this is also an ideal system for modern lattice simulations, so we also expect more lattice calculations in this family. We would like to note that the lattice calculations are becoming more and more precise, and the precision of their numerical evaluation can be much higher than in QCD sum rules approach (See Ref. [36] and two very recent talks given in Quarkonium 2014 [37, 38] for more information).

We also obtained the two decay constants $f_{-,3/2}$ and $f_{-,5/2}$, both of which are important input parameters when performing the dynamical study relevant to the D -wave charmed-strange mesons. With the running of the LHCb experiment, it is an exciting time to explore the higher charmed-strange mesons. The experimental and theoretical efforts will establish the charmed-strange meson family step by step, which is a research area full of challenges and opportunities.

Acknowledgments

This project is supported by the National Natural Science Foundation of China under Grants No. 11205011, No. 11475015, No. 11375024, No. 11222547, No. 11175073, No. 11035006, and NO. 11261130311, the Ministry of Education of China (SRFDP under Grant No. 20120211110002 and the Fundamental Research Funds for the Central Universities), and the Fok Ying-Tong Education Foundation (No. 131006).

-
- [1] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **90**, 242001 (2003).
 - [2] D. Besson *et al.* [CLEO Collaboration], Phys. Rev. D **68**, 032002 (2003) [Erratum-ibid. D **75**, 119908 (2007)].
 - [3] J. Brodzicka *et al.* [Belle Collaboration], Phys. Rev. Lett. **100**, 092001 (2008).
 - [4] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. D **80** (2009) 092003.
 - [5] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **97**, 222001 (2006).
 - [6] X. Liu, Int. J. Mod. Phys. Conf. Ser. **2**, 147 (2011).
 - [7] R. Aaij *et al.* [LHCb Collaboration], arXiv:1407.7574 [hep-ex].
 - [8] R. Aaij *et al.* [LHCb Collaboration], arXiv:1407.7712 [hep-ex].
 - [9] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
 - [10] Z. F. Sun and X. Liu, Phys. Rev. D **80**, 074037 (2009).
 - [11] Q. T. Song, D. Y. Chen, X. Liu and T. Matsuki, arXiv:1408.0471 [hep-ph].
 - [12] Z. G. Wang, arXiv:1408.6465 [hep-ph].
 - [13] S. Godfrey and K. Moats, arXiv:1409.0874 [hep-ph].
 - [14] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979).

- [15] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127**, 1 (1985).
- [16] B. Grinstein, Nucl. Phys. B **339**, 253 (1990).
- [17] E. Eichten and B. R. Hill, Phys. Lett. B **234**, 511 (1990).
- [18] A. F. Falk, H. Georgi, B. Grinstein and M. B. Wise, Nucl. Phys. B **343**, 1 (1990).
- [19] E. Bagan, P. Ball, V. M. Braun and H. G. Dosch, Phys. Lett. B **278**, 457 (1992).
- [20] M. Neubert, Phys. Rev. D **45**, 2451 (1992).
- [21] M. Neubert, Phys. Rept. **245**, 259 (1994).
- [22] D. J. Broadhurst and A. G. Grozin, Phys. Lett. B **274**, 421 (1992).
- [23] P. Ball and V. M. Braun, Phys. Rev. D **49**, 2472 (1994).
- [24] T. Huang and C. W. Luo, Phys. Rev. D **50**, 5775 (1994).
- [25] Y. B. Dai, C. S. Huang, M. Q. Huang and C. Liu, Phys. Lett. B **390**, 350 (1997).
- [26] Y. B. Dai, C. S. Huang and H. Y. Jin, Z. Phys. C **60**, 527 (1993).
- [27] Y. B. Dai, C. S. Huang and M. Q. Huang, Phys. Rev. D **55**, 5719 (1997).
- [28] P. Colangelo, F. De Fazio and N. Paver, Phys. Rev. D **58**, 116005 (1998).
- [29] Y. B. Dai, C. S. Huang, C. Liu and S. L. Zhu, Phys. Rev. D **68**, 114011 (2003).
- [30] P. Colangelo, G. Nardulli, A. A. Ovchinnikov and N. Paver, Phys. Lett. B **269**, 201 (1991).
- [31] P. Colangelo, G. Nardulli and N. Paver, Phys. Lett. B **293**, 207 (1992).
- [32] <http://www.feynncalc.org/>.
- [33] B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006).
- [34] Y. B. Dai and S. L. Zhu, Phys. Rev. D **58**, 074009 (1998).
- [35] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014).
- [36] C. B. Lang, L. Leskovec, D. Mohler, S. Prelovsek and R. M. Woloshyn, Phys. Rev. D **90**, 034510 (2014).
- [37] D. Mohler, talk given in Quarkonium 2014, 10-14 November 2014 at CERN, <http://indico.cern.ch/event/278195/session/0/contribution/29/material/slides/0.pdf>.
- [38] R. Lewis, talk given in Quarkonium 2014, 10-14 November 2014 at CERN, <http://indico.cern.ch/event/278195/session/0/contribution/30/material/slides/0.pdf>.